

Art therapy

Clinical Policy ID: CCP.1400

Recent review date: 8/2025

Next review date: 12/2026

Policy contains: Aging; art therapy; cancer; dementia; depression; psychosis; schizophrenia; sexual abuse.

FirstChoice VIP Care has developed clinical policies to assist with making coverage determinations. FirstChoice VIP Care's clinical policies are based on guidelines from established industry sources, such as the Centers for Medicare & Medicaid Services (CMS), state regulatory agencies, the American Medical Association (AMA), medical specialty professional societies, and peer-reviewed professional literature. These clinical policies along with other sources, such as plan benefits and state and federal laws and regulatory requirements, including any state- or plan-specific definition of "medically necessary," and the specific facts of the particular situation are considered, on a case by case basis, by FirstChoice VIP Care when making coverage determinations. In the event of conflict between this clinical policy and plan benefits and/or state or federal laws and/or regulatory requirements, the plan benefits and/or state and federal laws and/or regulatory requirements shall control. FirstChoice VIP Care's clinical policies are for informational purposes only and not intended as medical advice or to direct treatment. Physicians and other health care providers are solely responsible for the treatment decisions for their patients. FirstChoice VIP Care's clinical policies are reflective of evidence-based medicine at the time of review. As medical science evolves, FirstChoice VIP Care will update its clinical policies as necessary. FirstChoice VIP Care's clinical policies are not guarantees of payment.

Coverage policy

Art therapy is clinically proven and, therefore, may be medically necessary (National Comprehensive Cancer Network, 2024; National Institute for Health and Care Excellence, 2014, 2016a, 2016b; Regev, 2018):

- For members in a case management or disease management program with a behavioral health diagnosis, including dementia, depression, psychosis, or schizophrenia.
- For children with a history of sexual abuse.
- For members with chronic diseases including cancer.
- For members older than age 65 to support mental well-being and independence by developing or maintaining social participation and preventing loneliness and social isolation.

Limitations

Art therapy is investigational/not clinically proven and, therefore, not medically necessary for post-traumatic stress disorder, as its effectiveness has not been proven for this condition (National Institute for Health and Care Excellence, 2018).

Alternative covered services

Alternate treatments depend on diagnosis and stage, and include behavioral therapies (cognitive behavioral therapy, trauma-focused cognitive behavioral therapy, single or group psychoanalytic behavioral therapy, and family intervention); pharmacologic therapies (antidepressants and antipsychotics); and alternative leisure activities.

Background

Art therapy applies art media, the creative process, and the final artwork as a therapeutic and healing process (Art Therapy Credentials Board, 2021b). For individuals experiencing developmental, medical, educational, social, or psychological impairments, art therapy is used to improve or restore the patient's functioning and sense of well-being.

Art therapy is used in inpatient, outpatient, and community settings (Art Therapy Credentials Board, 2021a). Art therapy may benefit patients in multiple ways, including by providing a sense of satisfaction and accomplishment, important to those feeling frustrated by their medical conditions; by helping those who are emotionally or verbally withdrawn to express themselves; and by helping those with life-limiting diseases, such as cancer, to find an identity outside of their condition and to make meaning of their experiences.

Entry into the profession of art therapy requires a master's degree (Art Therapy Credentials Board, 2021a). Educational standards for approved art therapy master programs are established and overseen by the American Art Therapy Association. The Art Therapy Credentials Board ensures that the educational and professional standards for being a qualified art therapist are met and maintained. The Board confers and administers four levels of professional credentials to art therapy practitioners, including board certification (Art Therapy Credentials Board, 2021b).

Findings

Guidelines

The National Institute for Health and Care Excellence guidance recommends offering art-based therapies as part of a multidisciplinary package of care for adults and children with psychosis or schizophrenia, and as a way to promote social participation and mental well-being in older adults (National Institute for Health and Care Excellence, 2014, 2016a, 2016b). Conversely, its guideline on post-traumatic stress disorder does not recommend art therapy due to insufficient evidence (National Institute for Health and Care Excellence, 2018). A German psychiatric guideline lists art therapy among optional interventions for anxiety disorders but notes that the evidence remains limited (Bandelow, 2022). In oncology, the National Comprehensive Cancer Network recommends incorporating creative arts interventions, such as art and music, into psychosocial support for cancer-related distress (National Comprehensive Cancer Network, 2024).

Evidence reviews

Systematic reviews consistently report that art therapy has positive effects on psychological outcomes across varied populations but highlight heterogeneity, small sample sizes and methodological limitations. A 2018 systematic review (Regev, 2018) noted promising results for quality-of-life outcomes in cancer and mixed findings in other conditions. A large JAMA Network meta-analysis of 50 randomized trials ($n = 2,766$) found that art therapy improved only about 18% of reported outcomes; effect sizes were small (change-score standardized mean difference = 0.38 and post-test SMD = 0.19), and study quality was generally low (Joschko, 2024). More recent reviews echo these concerns, while also providing quantitative data for specific populations.

Cancer and other chronic diseases

In adults with cancer, a 2025 meta-analysis of 22 randomized controlled trials reported that art-based interventions moderately improved quality of life (SMD = 0.38, 95% CI [0.25, 0.51]) but did not significantly reduce fatigue overall; benefits were greater when interventions lasted at least two months and included music or dance components (Zou, 2025). Another 2024 review of mixed RCTs and quasi-experimental studies in patients receiving chemotherapy found that painting, drawing or sculpting improved overall and social quality-of-life measures but not depressive symptoms (Abu-Odah, 2024). Evidence in pediatric oncology is emerging; a 2025

meta-analysis of three RCTs and five quasi-experimental studies involving 452 children with cancer showed that art therapy significantly reduced anxiety and depression and qualitatively improved stress and anger (Zhou, 2025 a). Separate systematic reviews in stroke rehabilitation describe improvements in limb function, fine motor ability and emotional wellbeing from creative arts-based therapies (Lo, 2018; Liu, 2024), and a small meta-analysis of five trials in diabetes reported reductions in depression (SMD = -1.36, 95% CI [-1.63, -1.09]) and blood glucose without changes in anxiety or glycated hemoglobin (Yang, 2021).

Mental health conditions in children and adults

An updated 2025 systematic review and meta-analysis of art-based interventions for depression among children and adolescents included 12 randomized controlled trials and found that art therapy significantly reduced depressive symptoms (SMD = -0.72, $p = 0.01$) and that visual arts, music, and drama were all beneficial (Zhang, 2025). Group painting therapy for depressed adolescents has shown improvements in positive and negative emotion scores and in cognitive and executive functioning (Yuan, 2021). In schizophrenia, a 2024 meta-analysis of 31 RCTs covering painting, calligraphy and handcrafts found small-to-moderate improvements in positive symptoms (SMD = 0.41), moderate improvements in negative symptoms (SMD = 0.70) and depression (SMD = 0.61) and large improvements in anxiety (SMD = 0.91); longer weekly sessions and female participants were associated with greater benefit (Du, 2024). Evidence for eating disorders remains inconclusive (Pedra Cruz Bettin, 2024). A 2025 meta-analysis focusing on breast-cancer survivors' mental health reported that art therapy significantly reduced anxiety and depression, with effect sizes of -0.771 for depression and -1.594 for anxiety (Huang, 2025 a).

Cognition, dementia and aging

In a 2025 randomized controlled trial of 78 dyads, a creative expressive art-based storytelling program delivered twice weekly for 12 weeks reduced behavioral and psychological symptoms of dementia, caregiver distress and burden, and improved communication and quality of life for persons with dementia; these benefits persisted for up to 24 weeks (Zhuo, 2025 b). A mixed-methods cluster trial of a social-art intervention for older adults with mild cognitive impairment found significant improvement in global cognitive function immediately after a 14-week program ($\beta = 2.85$; 95% CI [1.27, 4.44]) but no sustained effects at six months and no significant changes in psychosocial indicators or functional abilities (Huang, 2025 b). Meta-analyses continue to show that art therapy reduces depressive symptoms in older adults (Jenabi, 2023) and improves quality of life, cognitive and emotional functions in persons with dementia (Emblad, 2021; de Souza, 2022). The 2024 scoping review of creative activities for stroke survivors (Liu, 2024) and a 2024 systematic review on mild cognitive impairment (Chiang, 2024) suggest potential cognitive and psychological benefits, but emphasize the need for long-term studies.

Developmental conditions and neurodiversity

A 2024 systematic review of art and music therapy for children with autism spectrum disorder and attention-deficit hyperactivity disorder concluded that creative arts therapies improve symptomatology, behavior, communication, and social, cognitive, and emotional skills, and participants reported high satisfaction (Martínez-Vérez, 2024). Evidence for the effectiveness of art therapy in addressing language disorders or learning disabilities remains limited.

Trauma and stress-related conditions

Evidence for the effectiveness of art therapy in trauma treatment is mixed. A 2015 review of six comparative studies in adults reported reductions in trauma symptoms and depression in some trials but concluded that methodological weaknesses preclude firm conclusions (Schouten, 2015). Subsequent uncontrolled studies of art therapy in refugee children and mixed pediatric populations suggest potential benefits but are insufficient for independent recommendations (Annous, 2022; Braitto, 2022). A 2018 review found that art therapy interventions reduced stress markers such as self-reported scales, cortisol, and blood pressure in most studies and helped

bereaved individuals find meaning and feel connected to loved ones (Martin, 2018; Weiskittle, 2018). A more recent randomized trial comparing narrative writing, art therapy, and control conditions in mothers of premature infants found no significant differences in stress reduction (Martínez-Shaw, 2023). Mandala art therapy may improve negative emotions and hope, relieve pain, and reduce physiological indicators of stress, but the evidence is uncertain (Zhang, 2024).

Post-traumatic stress disorder

While art therapy for post-traumatic stress disorder (PTSD) has attracted research interest, the current evidence base is insufficient to support its clinical use. A recent 2025 meta-analysis reported a significant reduction in PTSD symptoms (SMD = -1.98, 95% CI [-3.80, -0.16]) among 665 participants but noted extreme between-study heterogeneity ($I^2 \approx 98\%$) and a limited number of trials, substantially reducing confidence in these findings (Wang, 2025). This marked variability indicates that included studies may have examined fundamentally different interventions or populations, undermining the reliability of pooled effect estimates. Previous evidence is similarly inconclusive: a 2015 systematic review of six controlled trials found significant trauma symptom improvements in only half the studies, and just one reported decreased depression (Schouten, 2015). Small, uncontrolled studies with refugee children and other pediatric groups have described improvements in stress, hope, and coping after art-based interventions (Annous, 2022; Braito, 2022; Zhang, 2024), but these studies lack methodological rigor and generalizability due to small sample sizes, absence of control groups, and unstandardized outcome measures. Consequently, the National Institute for Health and Care Excellence (NICE) guideline on PTSD does not recommend art therapy for treatment, reflecting its investigational status until more robust, high-quality evidence becomes available (NICE, 2018; last reviewed 2024).

In 2025, we reorganized and condensed the findings section and added several new systematic reviews and meta-analysis. No policy changes were warranted.

References

On July 21, 2025, we searched PubMed and the databases of the Cochrane Library, the U.K. National Health Services Centre for Reviews and Dissemination, the Agency for Healthcare Research and Quality, and the Centers for Medicare & Medicaid Services. Search terms were “Art therapy” (MeSH), “creative therap*,” and “activity therap*.” We included the best available evidence according to established evidence hierarchies (typically systematic reviews, meta-analyses, and full economic analyses, where available) and professional guidelines based on such evidence and clinical expertise.

Abu-Odah H, Sheffield D, Hogan S, Yorke J, Molassiotis A. Effectiveness of creative arts therapy for adult patients with cancer: A systematic review and meta-analysis. *Support care cancer*. 2024;32(7):430. Doi: 10.1007/s00520-024-08582-4.

Annous N, Al-Hroub A, El Zein F. A systematic review of empirical evidence on art therapy with traumatized refugee children and youth. *Front Psychol*. 2022;13:811515. Doi: 10.3389/fpsyg.2022.811515.

Art Therapy Credentials Board, Inc. About the credentials. <https://atcb.org/about-the-credentials/>. Published 2021.(a)

Art Therapy Credentials Board, Inc. What is art therapy? <https://atcb.org/what-is-art-therapy/>. Published 2021.(b)

Bandelow B, Werner AM, Kopp I, Rudolf S, Wiltink J, Beutel ME. The German guidelines for the treatment of anxiety disorders: First revision. *Eur Arch Psychiatry Clin Neurosci*. 2022;272(4):571-582. Doi: 10.1007/s00406-021-01324-1.

Braito I, Rudd T, Buyuktasgin D, Ahmed M, Glancy C, Mulligan A. Review: Systematic review of effectiveness of art psychotherapy in children with mental health disorders. *Ir J Med Sci*. 2022;191(3):1369-1383. Doi: 10.1007/s11845-021-02688-y.

Chiang L, Cheong D, Cordato NJ, Smerdely P. Visual art therapy and its effects in older people with mild cognitive impairment: A systematic review. *Int J Geriatr Psychiatry*. 2024;39(1):e6053. Doi: 10.1002/gps.6053.

de Souza LBR, Gomes YC, de Moraes MGG. The impacts of visual art therapy for elderly with neurocognitive disorder: A systematic review. *Dement Neuropsychol*. 2022;16(1):8-18. Doi: 10.1590/1980-5764-dn-2021-0042.

Du SC, Li CY, Lo YY, et al. Effects of visual art therapy on positive symptoms, negative symptoms, and emotions in individuals with schizophrenia: A systematic review and meta-analysis. *Healthcare (Basel)*. 2024;12(11):1156. Doi: 10.3390/healthcare12111156.

Emblad SYM, Mukaetova-Ladinska EB. Creative art therapy as a non-pharmacological intervention for dementia: A systematic review. *J Alzheimers Dis Rep*. 2021;5(1):353-364. Doi: 10.3233/ADR-201002.

Huang T, Yin X. Exploring the benefits of art therapy for anxiety and depression in women with breast cancer: a meta-analytical systematic review. *J Psychiatr Res*. 2025 (a); Epub ahead of print. doi:10.1016/j.jpsychires.2025.05.027..

Huang C, Yan Y, Tam WWS, et al. Effects of an integrated social-art intervention on cognitive and psychosocial outcomes among older adults with mild cognitive impairment in nursing homes: a mixed methods study. *BMC Med*. 2025(b);23(1):256. Doi:10.1186/s12916-025-04085-z.

Jenabi E, Bashirian S, Ayubi E, Rafiee, Bashirian M. The effect of the art therapy interventions on depression symptoms among older adults: A meta-analysis of controlled clinical trials. *J Geriatr Psychiatry Neurol*. 2023;36(3):185-192. Doi: 10.1177/08919887221130264.

Joschko R, Klatte C, Grabowska WA, Roll S, Berghöfer A, Willich SN. Active visual art therapy and health outcomes: A systematic review and meta-analysis. *JAMA Netw Open*. 2024;7(9):e2428709. Doi:10.1001/jamanetworkopen.2024.28709.

Liu S, Huang X, Liu Y, Yue J, Li Y, Chen L. A scoping review of the use of creative activities in stroke rehabilitation. *Clin Rehabil*. 2024;38(4):497-509. Doi: 10.1177/02692155241227049.

Lo TLT, Lee JLC, Ho RTH. Creative arts-based therapies for stroke survivors: A qualitative systematic review. *Front Psychol*. 2018;9:1646. Doi: 10.3389/fpsyg.2018.01646.

Martin L, Oepen R, Bauer K, et al. Creative arts interventions for stress management and prevention – A systematic review. *Behav Sci (Basel)*. 2018;8(2):28. Doi: 10.3390/bs8020028.

Martínez-Shaw ML, Sánchez-Sandoval Y. Effective stress intervention programs for parents of premature children: A systematic review. *Stress Health*. 2023;39(2):236-254. Doi: 10.1002/smi.3194.

Martínez-Vérez V, Gil-Ruiz P, Domínguez-Lloria S. Interventions through art therapy and music therapy in autism spectrum disorder, adhd, language disorders, and learning disabilities in pediatric-aged children: A systematic review. *Children (Basel)*. 2024;11(6). Doi: 10.3390/children11060706.

National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®). Distress management. Version 1.2025. www.nccn.org. Published November 18, 2024.

National Institute for Health and Care Excellence. Mental wellbeing and independence for older people. Quality statement 3: Social participation. Quality standard 137. www.nice.org.uk/guidance/qs137. Published December, 2016.(a)

National Institute for Health and Care Excellence. Post-traumatic stress disorder. NICE guideline [NG116]. <https://www.nice.org.uk/guidance/ng116>. Published December 2018. Last reviewed April 8, 2025.

National Institute for Health and Care Excellence. Psychosis and schizophrenia in adults: Recognition and management. Clinical guideline 178. www.nice.org.uk/guidance/cg178. Published February 2014. Updated March 2014. Last Reviewed September 19, 2024.

National Institute for Health and Care Excellence. Psychosis and schizophrenia in children and young people: Recognition and management. Clinical guideline [CG155]. www.nice.org.uk/guidance/cg155. Published 2013. Updated October 2016 (b). Last reviewed September 19, 2024.

Pedra Cruz Bettin B, Urquiza Nogueira L, Bertasso de Araujo PA, Antunes LC. Visual art- and music-based interventions as adjuvants in the treatment of eating disorders: A systematic review and a theoretical model. *Arts & health*. 2024;16(2):167-188. Doi: 10.1080/17533015.2023.2218408.

Regev D, Cohen-Yatziv L. Effectiveness of art therapy with adult clients in 2018 — what progress has been made? *Front Psychol*. 2018;9:1531. Doi: 10.3389/fpsyg.2018.01531.

Schouten KA, de Niet GJ, Knipscheer JW, Kleber RJ, Hutschemaekers GJ. The effectiveness of art therapy in the treatment of traumatized adults: A systematic review on art therapy and trauma. *Trauma Violence Abuse*. 2015;16(2):220-228. Doi: 10.1177/1524838014555032.

Wang J, Zhang B, Yahaya R, Abdullah AB. Colors of the mind: a meta-analysis of creative arts therapy as an approach for post-traumatic stress disorder intervention. *BMC Psychol*. 2025;13(1):32. Doi:10.1186/s40359-025-02361-4.

Weiskittle RE, Gramling SE. The therapeutic effectiveness of using visual art modalities with the bereaved: A systematic review. *Psychol Res Behav Manag*. 2018;11:9-24. Doi: 10.2147/prbm.s131993.

Yang Q, Shao Q, Xu Q, Shi H, Li L. Art therapy alleviates the levels of depression and blood glucose in diabetic patients: A systematic review and meta-analysis. *Front Psychol*. 2021;12:639626. Doi: 10.3389/fpsyg.2021.639626.

Yuan Z, Li B, Ye X, Zhu X. Systematic review and meta-analysis of the effects of group painting therapy on the negative emotions of depressed adolescent patients. *Ann Palliat Med*. 2021;10(10):10744-10755. Doi: 10.21037/apm-21-2474.

Zhang B, Yang L, Sun W, Xu P, Ma H, Abdullah AB. The effect of the art therapy interventions to alleviate depression symptoms among children and adolescents: a systematic review and meta-analysis. *Clinics (Sao Paulo)*. 2025;80:100683. Doi:10.1016/j.clinsp.2025.100683

Zhang MQ, Liu X, Huang Y. Does mandala art improve psychological well-being in patients? A systematic review. *J Integr Complement Med*. 2024;30(1):25-36. Doi: 10.1089/jicm.2022.0780.

Zhou S, Li H, Yang Y, et al. Effects of art therapy on psychological outcomes among children and adolescents with cancer: a systematic review and meta-analysis. *BMC Complement Med Ther*. 2025(a);25(1):149. Doi:10.1186/s12906-025-04866-2.

Zhuo X, Yan Y, Lin R, et al. Effects of an art-based intervention in older adults with dementia: a randomized controlled trial [published correction appears in *Sci Rep*. 2025 (b);15(1):15303. Doi: 10.1038/s41598-025-99769-0.

Zou Y, Zhai S, Dong J, Zhao T, Yang J. The effect of art therapy on quality of life and fatigue in breast cancer patients: a meta-analysis. *Front Oncol*. 2025;15:1495887. Doi:10.3389/fonc.2025.1495887.

Policy updates

7/2018: initial review date and clinical policy effective date: 9/2018

8/2019: Policy references updated.

8/2020: Policy references updated.

8/2021: Policy references updated.

8/2022: Policy references updated.

8/2023: Policy references updated.

8/2024: Policy references updated.

8/2025: Policy references updated.